arXiv:1303.2334v2 [math.NT] 17 Mar 2013

Galois groups associated to generic Drinfeld modules and a conjecture of Abhyankar

Florian Breuer
Stellenbosch University, Stellenbosch, South Africa
fbreuer@sun.ac.za

March 21, 2013

Abstract

Let φ be a rank r Drinfeld $\mathbb{F}_q[T]$-module determined by $\varphi_T(X) = TX + g_1X^q + \cdots + g_{r-1}X^{q^{r-1}} + X^{q^r}$, where $g_1, \ldots, g_{r-1}$ are algebraically independent over $\mathbb{F}_q(T)$. Let $N \in \mathbb{F}_q[T]$ be a polynomial, and $k/\mathbb{F}_q$ an algebraic extension. We show that the Galois group of $\varphi_N(X)$ over $k(T, g_1, \ldots, g_{r-1})$ is isomorphic to $\mathrm{GL}_r(\mathbb{F}_q[T]/N\mathbb{F}_q[T])$, settling a conjecture of Abhyankar.

1 Introduction

Let $\mathbb{F}_q$ be the finite field of q elements, $k/\mathbb{F}_q$ an algebraic extension and let $T, g_1, g_2, \ldots, g_{r-1}$ be algebraically independent over $\mathbb{F}_q$, for some $r \geq 2$. Let $A := \mathbb{F}_q[T]$ be the polynomial ring over $\mathbb{F}_q$, $F := k(T)$ the rational function field over k and $K := k(T, g_1, \ldots, g_{r-1})$.

Consider the Drinfeld A-module φ of rank r defined over K by

$$\varphi_T(X) = TX + g_1X^q + \cdots + g_{r-1}X^{q^{r-1}} + X^{q^r}.$$

See [5, Chap. 4] for a background on Drinfeld modules. For any $N = a_0 + a_1T + \cdots + a_nT^n \in A$, the polynomial $\varphi_N(X) \in K[X]$ is constructed by iterating φ_T as follows:

$$\varphi_N(X) = a_0 + a_1\varphi_T(X) + a_2(\varphi_T \circ \varphi_T)(X) + \cdots + a_n \underbrace{(\varphi_T \circ \cdots \circ \varphi_T)}_{n \text{ times}}(X).$$

Denote by K_N the splitting field of $\varphi_N(X)$ over K. Equivalently, K_N is obtained by adjoining the elements of the N-torsion module $\varphi[N]$ of φ to K. Since $\varphi[N] \cong (A/NA)^r$ as A-modules, and $\mathrm{Gal}(K_N/K)$ commutes with the A-module structure, we see that $\mathrm{Gal}(K_N/K) \hookrightarrow \mathrm{GL}_r(A/NA)$ (the embedding depends on a choice of basis of $\varphi[N]$).

The goal of this note is to prove the following.

Theorem 1 *$\mathrm{Gal}(K_N/K) \cong \mathrm{GL}_r(A/NA)$.*

This is S.S. Abhyankar's "Generalized Iteration Conjecture", see [1, §19]. It is also the Drinfeld module analogue of the following theorem of Weber, see [8, Cor. 1, p68]. Let E be an elliptic curve with transcendental j-invariant j, defined over $\mathbb{Q}(j)$, and $n > 1$, then $\mathrm{Gal}(\mathbb{Q}(j, E[n])/\mathbb{Q}(j)) \cong \mathrm{GL}_2(\mathbb{Z}/n\mathbb{Z})$.

When $N = T$, Theorem 1 can be traced back to E.H. Moore [9]:

Theorem 2 (Moore) $\mathrm{Gal}(K_T/K) \cong \mathrm{GL}_r(A/TA) = \mathrm{GL}_r(\mathbb{F}_q)$.

Proof. See [2, §3] for a particularly simple proof. $\square$

A number of other special cases of Theorem 1 are known, for example the case $r = 1$ is due to Carlitz [3], the case $r = 2$ is closely related to a result of Joshi [7], and the case $N = T^n$ with $r \geq 1$ arbitrary was proved independently by Thiery [12] and by Abhyankar and Sundaram [2]. Theorem 1 has also been proved under a variety of hypotheses on $N \in A$ by Abhyankar and his students, see [1, §19] and the references therein.

2 Relative Galois groups

Let $N, M \in A$ and define

$$\begin{aligned} G(MN, M) &:= \{g \in \mathrm{GL}_r(A/MNA) \mid g \equiv 1 \bmod M\} \\ &= \ker\big(\mathrm{GL}_r(A/MNA) \longrightarrow \mathrm{GL}_r(A/MA)\big). \end{aligned}$$

Our approach will be to show the following.

Theorem 3 $\mathrm{Gal}(K_{MN}/K_M) \cong G(MN, M)$.

Clearly, Theorem 3 follows from Theorem 1. Conversely, we show that Theorem 1 follows if Theorem 3 holds for $M = T$.

Proof of Theorem 1.

$$\begin{tikzcd} & K_{NT} \arrow[dl, dash, "G(NT{,}T)"'] \arrow[dr, dash, "G(NT{,}N)"] & \\ K_T \arrow[dr, dash, "\mathrm{GL}_r(A/TA)"'] & & K_N \arrow[dl, dash] \\ & K & \end{tikzcd}$$

Suppose that $\mathrm{Gal}(K_{NT}/K_T) \cong G(NT, T)$. Consider the field extensions in the diagram. We have $\mathrm{Gal}(K_T/K) \cong \mathrm{GL}_r(A/TA)$ by Theorem 2, thus $[K_{NT} : K] = \#G(NT, T) \cdot \#\,\mathrm{GL}_r(A/TA) = \#\,\mathrm{GL}_r(A/NTA)$ and it follows that $\mathrm{Gal}(K_{NT}/K) \cong \mathrm{GL}_r(A/NTA)$. Now from the action of $\mathrm{Gal}(K_{NT}/K)$ on $\varphi[N]$ we see that $\mathrm{Gal}(K_{NT}/K_N) \cong G(NT, N)$ and finally $\mathrm{Gal}(K_N/K) \cong \mathrm{GL}_r(A/NTA)/G(NT, N) \cong \mathrm{GL}_r(A/NA)$. $\square$

3 Drinfeld moduli schemes

The key point is to recognize the groups $G(MN, M)$ as the Galois groups of étale morphisms between Drinfeld moduli schemes, so we introduce these next. A useful reference is the careful article of Hubschmid [6].

Let $N \in A$ be non-constant, and consider the functor $\mathcal{F}_N^r : F-\text{Schemes} \to \text{Sets}$, which maps an F-scheme S to the set of isomorphism classes of rank r Drinfeld A-modules with level-N structure over S. Then $\mathcal{F}_N^r$ is representable by a non-singular affine scheme M_N^r over $\mathrm{Spec}(F)$, called a Drinfeld moduli scheme; see [6, §2] for details.

Theorem 4 (Drinfeld) *Let $M, N \in A$ be non-constant. Then the natural projection $M_{MN}^r \to M_M^r$ is an étale Galois cover over F with group $G(MN, M)$.*

Proof. The ideas are due to Drinfeld [4], see [6, Prop. 3.1.3] for a detailed proof. $\square$

Next, we need the following result.

Theorem 5 *Let $N \in A$ be non-constant. Then M_N^r is irreducible as a scheme over* $\mathrm{Spec}(F)$.

Proof. This is [6, Cor. 3.4.5] in the case where $k = \mathbb{F}_q$. In our situation, the same proof still works, namely there exists a morphism $M_N^r \to M_N^1$ defined over F by [6, Cor. 3.4.4], whose geometric fibres are precisely the geometrically irreducible components of M_N^r by [6, Prop. 3.4.2]. It remains to check that M_N^1 is irreducible. For this we recall [4, Thm 1] that $M_N^1 = \mathrm{Spec}(R_N^1 \otimes_A F)$ where R_N^1 is the integral closure of A in a certain class field over $\mathbb{F}_q(T)$ which splits completely above the place of $\mathbb{F}_q(T)$ with uniformizer $\frac{1}{T}$. As a result, this class field is a purely geometric extension of $\mathbb{F}_q(T)$, whereas F is a constant field extension of $\mathbb{F}_q(T)$, and so $R_N^1 \otimes_A F$ is a field, and M_N^1 is irreducible. $\square$

4 The case $N = T$

When $N = T$, the moduli scheme M_T^r has a nice explicit description, due to Pink [10, 11], since a Drinfeld A-module is uniquely determined by its level-T structure.

Let $V = \mathbb{F}_q^r$, $S_V = F[v \mid v \in V]$ the symmetric algebra of V over F, and $R_V = F[v, \frac{1}{v} \mid v \in V \smallsetminus \{0\}]$ the localization of S_V at $\prod_{v \in V \smallsetminus \{0\}} v$. We turn R_V into a graded ring by defining $\deg(v) = 1$ and $\deg(\frac{1}{v}) = -1$ for all $v \in V \smallsetminus \{0\}$, and $\deg(x) = 0$ for all $0 \neq x \in F$. Let $R_{V,0}$ denote the degree 0 component of R_V, then from [10, §7] we obtain the following result.

Theorem 6 (Pink) $M_T^r = \mathrm{Spec}(R_{V,0})$.

Moreover, the Drinfeld module φ endowed with a level-T structure $\lambda : V \xrightarrow{\sim} \varphi[T]$ defines a K_T-valued point of M_T^r, denoted $\eta_T : \mathrm{Spec}(K_T) \to M_T^r$, corresponding to

$$\eta_T^\sharp : R_{V,0} \longrightarrow K_T, \quad \text{induced by} \quad \frac{v_1}{v_2} \longmapsto \frac{\lambda(v_1)}{\lambda(v_2)}, \quad \forall v_1, v_2 \in V \smallsetminus \{0\}.$$

Since $\eta_T^\sharp$ is injective, η_T maps onto the generic point of M_T^r.

By Theorem 5, M_{NT}^r is irreducible, hence has a unique generic point. Now φ endowed with a level-NT structure defines a K_{NT}-valued point η_{NT} of M_{NT}^r, which lies above η_T, and hence also maps onto the generic point of M_{NT}^r.

Denote by $K_{T,0} \subset K_T$ the quotient field of the image of $\eta_T^\sharp$, then $K_T = K_{T,0}(\lambda(v))$ for any $0 \neq v \in V$, so K_T is a purely transcendental extension of $K_{T,0}$. Similarly, we denote by $K_{NT,0} \subset K_{NT}$ the quotient field of the image of $\eta_{NT}^\sharp$. It follows from Theorem 4 that $\mathrm{Gal}(K_{NT,0}/K_{T,0}) \cong G(NT,T)$. Therefore, the compositum $K_{NT,0} \cdot K_T$ in K_{NT} is Galois over K_T with group $G(NT,T)$, so $[K_{NT} : K_T] \geq \# G(NT,T)$. Since $\mathrm{Gal}(K_{NT}/K_T) \hookrightarrow G(NT,T)$ it follows that $\mathrm{Gal}(K_{NT}/K_T) \cong G(NT,T)$. This proves Theorem 3 for the case $M = T$, and so Theorem 1 follows. $\square$

Acknowledgement. I wish to thank the anonymous referee who read an earlier draft of this paper, in which a more complicated proof of Theorem 1 was proposed, and who outlined the proof presented here. This work was supported by NRF grant number BS2008100900027.

References

[1] S.S. Abhyankar, Resolution of singularities and modular Galois theory. *Bull. Amer. Math. Soc. (N.S.)* **38** (2001), no. 2, 131–169.

[2] S.S. Abhyankar and G.S. Sundaram, Galois theory of Moore-Carlitz-Drinfeld modules. *C. R. Acad. Sci. Paris Sér. I Math.* **325** (1997), no. 4, 349–353.

[3] L. Carlitz, A class of polynomials. *Trans. Amer. Math. Soc.* **43** (1938), no. 2, 167–182.

[4] V. L. Drinfeld, Elliptic modules (Russian), *Math. Sbornik*, **94** (1974), 594-627. Translated in *Math. USSR. S.*, **23** (1974), 561– 592.

[5] D. Goss, Basic structures in function field arithmetic, Springer-Verlag, 1996.

[6] P. Hubschmid, The André-Oort conjecture for Drinfeld modular varieties, to appear in *Compos. Math.*; arXiv:1201.5556v1[math.NT]

[7] K. Joshi, A family of étale coverings of the affine line, *J. Number Theory* **59** (1996), 414–418.

[8] S. Lang, Elliptic Functions, 2nd edition, *Graduate Texts in Mathematics* **112**, Springer-Verlag, 1987.

[9] E.H. Moore, A two-fold generalization of Fermat's theorem. *Bull. Amer. Math. Soc.* **2** (1896), no. 7, 189–199.

[10] R. Pink, Compactification of Drinfeld modular varieties and Drinfeld Modular Forms of Arbitrary Rank, to appear in *Manuscripta Math.*; arXiv:1008.0013v4[math.AG]

[11] R. Pink, S. Schieder, Compactification of a Drinfeld Period Domain over a Finite Field, to appear in *J. Algebraic Geometry*; arXiv:1007.4796v3[math.AG]

[12] A. Thiery, $\mathbb{F}_q$-linear Galois theory, *J. London Math. Soc.* (2) **53** (1996), 441–454.